

Please ensure that your abstract fits into one column on one page and complies with the *Instructions to Authors* available from the Abstract Submission web page.

CORE-MANTLE PARTITIONING OF VOLATILE
ELEMENTS AND THE ORIGIN OF VOLATILE
ELEMENTS IN EARTH AND MOON

K. RIGHTER¹, K. PANDO², L. DANIELSON², K.
NICKODEM³

¹ Mailcode KT, NASA Johnson Space Center, 2101
NASA Pkwy., Houston, TX 77058

² Jacobs JETS, NASA Johnson Space Center, 2101
NASA Pkwy., Houston, TX 77058

³ Dept. Geography, Syracuse University, 144 Eggers
Hall, Syracuse, NY 13244-1020

Depletions of volatile siderophile elements (VSE; Ga, Ge, In, As, Sb, Sn, Bi, Zn, Cu, Cd) in mantles of Earth and Moon, constrain the origin of volatile elements in these bodies, and the overall depletion of volatile elements in Moon relative to Earth. A satisfactory explanation has remained elusive [1,2]. We examine the depletions of VSE in Earth and Moon and quantify the amount of depletion due to core formation and volatility of potential building blocks.

We calculate the composition of the Earth's PUM during continuous accretion scenarios with constant and variable fO_2 . Results suggest that the VSE can be explained by a rather simple scenario of continuous accretion leading to a high PT metal-silicate equilibrium scenario that establishes the siderophile element content of Earth's PUM near the end of accretion [3]. Core formation models for the Moon explain most VSE, but calculated contents of In, Sn, and Zn (all with $T_c < 750$ K) are all still too high after core formation, and must therefore require an additional process to explain the depletions in the lunar mantle. We discuss possible processes including magmatic degassing, evaporation, condensation, and vapor-liquid fractionation in the lunar disk.

References: [1] Ringwood, A.E. and Kesson, S. (1977) *The Moon* 16, 425-464. [2] Delano, J.W. and Ringwood, A.E. (1978) *Moon and the Planets* 18, 385-425. [3] Righter, K. (2011) *EPSL*.